


MSU Extension Publication Archive

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Corn Rootworm
Michigan State University
Cooperative Extension Service
Robert F. Ruppel, Entomology Department
March 1973
4 pages

The PDF file was provided courtesy of the Michigan State University Library

Scroll down to view the publication.



Corn Rootworm

By Robert F. Ruppel, Department of Entomology

The northern corn rootworm (Figure 1) has long been a pest of corn in certain Michigan fields. The western corn rootworm (Figure 2), a very similar pest, has recently invaded the state and is now established in our southwestern area. Both of these insects can damage only corn that is planted without rotation; a practice that is increasing in the state. The northern corn rootworm is definitely resistant to the chlorinated hydrocarbon insecticides (such as aldrin and chlordane) in some parts of Michigan. There is suspicion that the western corn rootworm, which has entered the state, is similarly resistant. The appearance of the western, the increasing acreage of constant corn, and the resistance of the worms to certain insecticides strongly indicate increasing problems with these pests in the near future. We must stay alert to avoid damage from these pests to our corn.

BIOLOGY

The western and northern corn rootworms are very similar in biology. Adult beetles of the rootworms emerge from the soil in late July and August. They are very active and hard to catch. They can be distinguished from nearly all other hard-shelled beetles found in flowering corn by their long antennae (feelers; see Figure 2). The beetles feed on the silks, tassels, soft kernels at the tips of the ears, and leaves of corn. (The western corn rootworm especially feeds on the leaves.) Their feeding on the early silks can result in a poor seed set. They lay their oval, yellowish eggs in the soil near the bases of corn plants. The adults are abundant in corn while it is flowering, and feed on the pollen of a number of plants when the corn matures. They are active in the field until the first hard frosts of the fall.

Figure 1. Northern corn rootworm adults feeding on a corn tassel. They are uniform yellow to green in color.



Figure 2. The western corn rootworm adult is yellow with black stripes down its wings. Both the northern and western corn rootworm adults have long antennae (feelers).

The eggs overwinter in the field and hatch when the soil warms up in the late spring. The larvae (rootworms) that hatch from these eggs are slender, cylindrical, white in color, and have a tan head and six small legs just in back of the head. The larvae vary in size to 1/2 inch when fully grown. The larvae tunnel and prune the roots of corn and can feed only on corn. Their feeding can destroy the roots (Figure 3) and reduces the vigor of the plant. More importantly, the damaged roots cause the stalks to lodge and pull out of the ground at harvest time. The lodging caused by the rootworm starts at soil level and the plants are commonly curved (goose-necked) at their bases (Figure 4). The greatest damage done by the rootworms is by the lodging and weakened roots that prevent mechanical harvesting of the crop. The larvae transform to a quiet stage (the pupae) in the soil in mid-to late-July and the adults emerge from these pupae. They have only one generation per year.

DETECTION

The damage of the rootworms occurs to the roots in the soil and is too often not noted until the corn lodges in late summer. This is after the damage has been done to that crop. The damage of the rootworms can be anticipated in time to prevent its occurrence if you will **Check Your Fields** for the pest.

Corn rootworms can be a problem only where corn follows corn. Reports of damage have been frequent after three years or more of constant corn, with few reports of damage to second-year corn. Damage has been more common on the "prairie" soils in the southern and eastern fringes of the state, but scattered fields with rootworms occur throughout the southern third of the state. Check for rootworms in any corn grown without rotation. Be especially careful if the field has been in corn for more than two years, and even more so if rootworms are prevalent in your area.

To check corn for rootworms:

(a) Examine tassels and silks for the presence of adults at flowering. This is the best time to detect the pest and prepare for its control during the next season. About one adult per plant indicates a rootworm problem the following year.

(b) Examine lodged and goose-necked plants when plants are mature for evidence of root tunneling. Smaller roots may be completely rotted off by this time, but brace roots may still show tunnels. The crop will need protection the following year if plants with goose-necking and tunneled roots are common.

(c) Examine plants that are stunted or wilted in late June or early July or are lodged later in the season for larvae and tunnels of the rootworms. A cultivation



Figure 3. Roots on the right have been severely damaged by corn rootworm larvae; the roots on the left were protected from damage by an insecticide applied at planting time.

application of insecticides (described below) may be made to protect the crop if worms are found in time. Carefully recheck the field for adults at flowering time if rootworm damage is suspected.

If you are still unsure of the need to protect against corn rootworm, use this simple test:

(a) Treat two or three separate swaths of four to eight rows wide across the field with an insecticide (see below) if you suspect that you may need to control the rootworm. Compare the treated rows for larvae, lodging, and harvest problems with the adjacent, untreated corn.

(b) Leave similar swaths untreated and compare them with the adjacent treated rows if you are now using an insecticide for rootworm control and suspect that you do not need it.

CONTROL

Rotation—rootworms can only be a pest where corn follows corn without rotation. The best way to control the rootworm is to rotate the corn with any other crop. Infested fields should be placed in rotation, unless there are real benefits from growing corn-after-corn in that particular field. The advantages of growing corn without rotation should be reviewed when problems develop with the corn rootworms.



Figure 4. Lodging of corn caused by destruction of the roots by corn rootworm larvae. The stalks tilt right from soil level and often curve up (goose-neck) near their bases.

Insecticides

Insecticides should be used to control rootworms in infested fields where rotation is not feasible. Remember that insecticides are poisonous; handle, store, and apply them with great care. The label on the insecticide container has full instructions for safe, effective use of that specific insecticide. **READ THE LABEL** before buying any insecticide. Insecticides used to control rootworms are applied to the soil. They should be covered with soil immediately after application to prevent their exposure to high soil surface temperatures and to people, livestock, and wildlife. The equipment used to apply the insecticides should be carefully calibrated to deliver the exact amount of insecticide desired. Insecticides recommended for use in 1973 are given in the table. Check with your county agricultural extension agent for current recommendations in future years.

Pre-plant broadcast applications are made by spraying the liquid insecticide or applying a granular insecticide to the entire field prior to planting. A good practice is to apply the insecticide immediately before the final disking. Broadcast applications require more insecticides, and are, therefore, more expensive than band applications.

Planting band applications are made by spraying a liquid insecticide or applying a granular insecticide in a seven-inch band centered over the seed row. The band application may be made at planting or between the time of planting and seed germination. Apply the insecticide above, and out of contact with, the seed. Place the insecticide nozzle or spout between the seed spout and covering wheel of the planter. Make sure that the insecticide is covered with soil. A modified covering wheel or a light harrow in back of the insecticide applicator can be used. The planting band application is the least expensive and most convenient method of rootworm control on most farms.

Cultivation band applications are made by spraying a liquid insecticide or applying a granular insecticide in 3- to 4-inch bands on both sides of rows of growing corn. This requires special equipment, but is the only practical way to apply insecticides for rootworm control of growing plants. A cultivator shoe should be used to cover the insecticide immediately after application.

Some Simple Rules—

- Check Your Fields for Corn Rootworms
- Rotate Crops in Infested Fields Whenever Feasible
- Use Insecticides Safely When They are Needed
- Cover the Insecticide with Soil Immediately after Application

Insecticides Recommended for Corn Rootworm Control in 1973^a

Type Application	Insecticide	Lbs. Active Insecticide/Acre	Notes
Pre-plant broadcast	diazinon	4	
	chlordane ^b	4	Non-dairy farms only
	BUX	4	
Planting band	diazinon	1 1/2 ^c	
	chlordane ^b	2 ^c	Non-dairy farms only
	carbofuran (Furadan)	3/4 ^c	Hazardous
	phorate (Thimet)	1 ^c	Hazardous
	BUX	1 ^c	
	Dasanit	1 ^c	Hazardous
	Dyfonate	1 ^c	Hazardous
	profos (Mocap, Jolt)	1 ^c	
Cultivation band	phorate (Thimet)	1 ^c	Hazardous; apply in June
	BUX	1 ^c	Apply 30-40 days after planting or when plants are 8-10 inches tall.
	diazinon	1 ^c	
	Dyfonate	1 ^c	Hazardous
	profos (Mocap, Jolt)	1 ^c	
	Dasanit	1 ^c	Hazardous
	carbofuran (Furadan)	3/4 ^c	Hazardous

^a Check with your County Agricultural Extension Agent for insecticides recommended in future years.

^b Do not use chlordane where other chlorinated hydrocarbon insecticides (such as aldrin and heptachlor) have failed to control the rootworms in previous years.

^c The amounts given in band applications are for use with 40 inch row widths. Adjust this amount when other row widths are used by the equation:

adjusted amount = $\frac{40 A}{W}$; where A is the amount of active insecticide/A given in the table and W is the row width, in inches, that will be used. For example, if a 32 inch row width is to be used with an insecticide listed at 1 lb. per acre:

$$\text{adjusted amount} = \frac{40 \times 1}{32} = 1 \frac{1}{4} \text{ lb.}$$