

MSU Extension Publication Archive

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Verticillium Wilt of Ornamentals
Michigan State University
Cooperative Extension Service
Franklin Laemmlen
Department of Botany and Plant Pathology
April 1975
2 pages

The PDF file was provided courtesy of the Michigan State University Library

Scroll down to view the publication.

VERTICILLIUM WILT OF ORNAMENTALS

NO. 32

EXTENSION BULLETIN E-838

APRIL 1975

BY FRANKLIN LAEMMLEN

Department of Botany and Plant Pathology

What is it?

Verticillium wilt is a vascular disease which attacks a wide variety of trees, shrubs and herbaceous ornamentals. The disease is caused by a fungus, *Verticillium albo-atrum*, which lives in the soil and enters the plant through the roots. In the plant, the fungus spreads upward, mainly in the wood, disrupting water movement and normal plant functions.

Symptoms

The disease may appear in mild, chronic or lethal form. Poor vigor and sparse growth may be the only symptoms (Fig. 1). Yellowing, marginal and interveinal browning, wilting and/or death of leaves may occur. The affected plant typically shows symptoms of mild to severe water stress. These often appear in one branch or one

side or sector of the plant (Fig. 2). In its lethal form Verticillium will cause a sudden and total collapse of the plant. The wood of affected plants shows discoloration in the form of streaks or bands (Fig. 3). These streaks may be scattered throughout the wood if the plant is chronically infected or may be confined to new sapwood in a new attack (Fig. 4). The color of the streaks can be from grayish to olive green (maples), to brown or near black in locust and elm. Streaks should be looked for at the base of the wilted branch or twig or in the trunk or roots on the affected side of the plant.

The above symptoms are not conclusive evidence of Verticillium wilt, as other diseases can cause similar symptoms. Positive identification can be made only by laboratory tests. Contact your County Cooperative Extension Service Office if you wish such tests to be made.



Fig. 1. (Left) Tree with a chronic Verticillium infection. The plant may persist in this condition for several years. Fig. 2. (Center) Note defoliated branch and one-sided flagging (arrows) of Verticillium affected tree. Fig. 3. (Right) Complete ring in current year's growth indicates why this branch died. Note initial Verticillium infection occurred 2 years earlier.



Fig. 4. Scattered streaks indicate a chronic infection, while streaks confined to the current year's growth indicate a new attack.

Trees and shrubs known to be susceptible to *Verticillium* wilt:

Ash, (Black, Blue European, Green and White)
Azalea (*Rhododendron molle*)
Barberry, Japanese
Boxwood, Korean
Catalpa, Western, Northern and Southern
Cherry
Elm, American and varieties Augustine Ascending, Henry Field, Littleford, and Moline Chinese (*Ulmus parvifolia*)
English Slippery
Goldenrain tree
Grapes
Horsechestnut
Kentucky coffee tree
Lilac
Linden, American and Little leaf
Locust, Black
Magnolia, Saucet and Star
Maple, Amur, Black, Hedge, Norway, and varieties Crimson King and Schwedleri, Red, Silver and Sugar
Oak, Pin and Red
Peach
Pear
Plum
Privet, Amut
Redbud
Rose daphne
Rose, Multiflora (most cultivated roses)
Russian olive
Sassafras
Smoketree
Sour gum
Spirea
Sumac, Fragrant, Smooth and Staghorn
Tree-of-heaven
Tulip tree
Viburnum

Replacement Species

Arborvitae
Beech
Birch
Fir
Flowering crabapple
Ginkgo
Hackberry
Hawthorn
Hickory
Holly
Honey locust
Hop hornbeam
Hornbeam
Juniper
Larch
Mountain ash
Mulberry
Oak, white and burr
Pawpaw (Asimina)
Pine
Poplar
Pyracantha (firethorn)
Serviceberry
Spruce
Sweet gum
Sycamore
Willow
Zelkova

Care of infected plants

There is no chemical cure available for *Verticillium* wilt. Trees showing general or severe wilt cannot be saved and should be cut down and disposed of by burning or removal to a dumping area. Infected wood **should not be chipped** (unless it is thoroughly heated in a compost pile manner) as the fungus can live for some time in chips and can grow as a saprophyte in the soil if adequate organic matter is present. In this manner, chip mulch could spread the disease.

Mild and chronically affected trees should be watered regularly to prevent water stress. Nitrogen fertilization to stimulate vigorous growth is also recommended. Prune out and destroy weak and dead branches.

Future plant selection

Where *Verticillium* wilt is found or suspected, resistant or tolerant species should be used in replacement plantings. If susceptible species must be used, soil fumigation by a licensed professional is required.

Verticillium wilt is not known to occur in yews and conifers. Also, a number of broadleaf species are not known to be susceptible to this disease. The species listed in the adjoining left column should be considered as replacement species for plants killed by *Verticillium* wilt.

Issued in furtherance of cooperative extension work in agriculture and home economics, acts of May 8, and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Gordon E. Guyer, Director, Cooperative Extension Service, Michigan State University, East Lansing, Michigan 48824.

1P-12M-4:75-UP