

## **MSU Extension Publication Archive**

Archive copy of publication, do not use for current recommendations. Up-to-date information about many topics can be obtained from your local Extension office.

Protein-Mineral Supplements for Growing and Finishing Cattle Fed Corn, Corn Silage Rations

Michigan State University

Cooperative Extension Service

Danny G. Fox, Beef Specialist, Department of animal Husbandry


J. Roy Black, Extension Economist, Department of Agricultural Economics

July 1976


4 pages

The PDF file was provided courtesy of the Michigan State University Library

**Scroll down to view the publication.**



# MICHIGAN BEEF PRODUCTION



Cooperative Extension Service  Michigan State University

## Protein-Mineral Supplements for Growing and Finishing Cattle Fed Corn, Corn Silage Rations

Danny G. Fox, Beef Specialist, Department of Animal Husbandry and  
J. Roy Black, Extension Economist, Department of Agricultural Economics

This fact sheet outlines the amount/head/day of protein-mineral supplement needed for growing and finishing cattle fed shelled corn, corn silage rations. Amounts are given for different weights, sexes, frame sizes, amounts of corn in the ration, and whether or not the corn silage was treated with a non-protein nitrogen (NPN) product. Also, the desired nutrient content of the protein-mineral supplements are given as well as the amounts of ingredients/1000 lb. batch soybean meal and urea based supplements. The supplementation programs are based upon Fact Sheet 1097, "Summary of Nutrient Requirements for Growing and Finishing Cattle." This fact sheet has two parts. The first describes the protein and mineral requirements of growing and finishing cattle, the average nutrient composition of corn, corn silage and supplements used to develop the feeding guidelines, and adjustment for sex and frame size of cattle. The second part contains a series of supplement feeding guidelines for different silage treatments.

### Summary of Nutrient Requirements Used

The protein and mineral requirements are based on the concept of *equivalent* weight. All weight groupings are referenced to an average framed steer, one that will reach a fatness of low choice at a weight of 1050 to 1100 lbs. The term equivalent weight is used to indicate weights at which animals of different frame sizes are equivalent in body composition, hence nutrient requirements. Small framed steers are defined as those reaching a fatness of low choice at 800 to 880 lbs., while large framed steers reach that point at

1200 to 1300 lbs. Small framed heifers reach low choice at 660 to 720 lbs., average framed at 800 to 880 lbs., and large framed at 940 to 1030 lbs.

Table 1 summarizes the protein requirements, as a % of ration dry matter, by sex, frame size, weight, and lbs. corn/head/day/cwt. body weight. Allowances were made for protein quality differences between corn grain and corn silage. As gain/day increases, as noted in Fact Sheet 1041, "Protein Requirements for Growing and Finishing Cattle," the *biologically available* protein required/day for tissue growth increases. Thus the biologically available protein required, as a % of ration dry matter, increases. But, the protein quality of corn grain is superior to that of corn silage; as a result, less total (crude) protein is required/lb. of weight gain on high grain than on high silage rations. The net effect is a small increase in total protein required as the corn/head/day fed increases for cattle with equivalent weights of less than 900 lbs. For equivalent weights of 900 lbs. to market weight, total protein required decreases as the corn in the ration increases since the major portion of the protein required at that stage of growth is for maintenance, not tissue gain. The mineral requirements for calcium, phosphorus, potassium, and salt are given in Table 2.

Table 3 summarizes the feeds used in developing the supplements and their nutrient values. Shelled corn is the primary energy feed. Protein feeds are 44% crude protein soybean meal and feed grade urea. Roughages include untreated corn silage and corn silage treated with sufficient NPN to raise its crude protein from 8% to 13% of dry matter. Minerals in-



**Table 2: Mineral Allowances**

Mineral	Animal, sex, frame size and weight (lbs)					
	<b>Heifer:</b>					
	Small frame	To 385	385-450	450-510	510-575	575-Market
	Average frame	To 480	480-560	560-640	640-720	720-Market
	Large frame	To 570	570-665	665-760	760-860	860-Market
<b>Steer:</b>						
	Small frame	To 480	480-560	560-640	640-720	720-Market
	Average frame	To 600	600-700	700-800	800-900	900-Market
	Large frame	To 720	720-840	840-960	960-1020	1020-Market
Mineral	Mineral, % of ration dry matter					
Calcium		.46	.40	.34	.28	.28
Phosphorus		.34	.28	.26	.24	.24
Potassium		.60	.60	.60	.60	.60
Salt		.25	.25	.25	.25	.25

**Table 3: Feed Nutrient Values (Michigan)**

Feed Description	Dry Matter	TDN	NE <sub>m</sub>	NE <sub>g</sub>	Total Protein	Calcium	Phosphorus	Potassium	Salt
	%	% of DM	Mcal/lb. DM	Mcal/lb. DM	% of DM	% of DM	% of DM	% of DM	% of DM
<b>Concentrates:</b>									
Corn	85	91	1.02	.67	10.0	.03	.40	.46	—
Soy 44	90	81	.88	.59	50.8	.36	.75	2.21	—
Urea	100	—	—	—	281.0	—	—	—	—
<b>Minerals:</b>									
Dicalcium Phosphate	100	—	—	—	—	23.0	18.0	—	—
Limestone	100	—	—	—	—	38.0	—	—	—
Calcium sulfate	100	—	—	—	—	20.3	—	—	—
Salt	100	—	—	—	—	—	—	—	100
<b>Forages:</b>									
Corn silage	32	70	.71	.45	8.0	.20	.21	.95	—
Corn silage with NPN added	32	70	.71	.45	13.0	.20	.21	.95	—

tion Evaluation Worksheet," to see if it will work your rations.

**Adjusting for Age, Sex, and Frame Size**

Table 4 gives the adjustment factor for amount of supplement fed/head/day for age, sex, and frame size. All amounts/head/day in subsequent sections are given for an average frame steer started on feed as a calf. But, a 780 lb. large framed steer has *equivalent* nutrient requirements, as a % of ration dry matter, to an average framed steer weighing 650 lbs. Thus pro-

tein-mineral supplement, as a % of ration dry matter, is the same for the 650 lb. average framed steer and the 780 lb. large framed steer. At 780 lbs., however, the large framed steer eats 13% more dry matter/day than the average framed steer and thus would need more lbs. of supplement/day.

*Example:*

A cattleman is starting a set of large framed heifer calves on an all corn silage ration; 2.1 lbs. of protein-mineral is recommended/head/day for an average framed steer started on feed as a calf. How should he

**Table 4: Adjustment Factors for the Protein-Mineral Supplement Feeding Rate for Frame Size, Sex, and Age.**

Age	Sex	Frame	Adjustment Factor, % of Average frame steer calf feeding rate
Calf	Heifer	Small	72
		Average	84
		Large	96
	Steer	Small	84
		Average	100
		Large	113
Yearling	Heifer	Small	80
		Average	93
		Large	105
	Steer	Small	93
		Average	110
		Large	124

adjust the daily feeding schedule? He should multiply the amount/head/day by .96 since the adjustment factor is 96%. That would yield,

$$2.1 \text{ lbs./hd./day} \times .96 = 2.0 \text{ lbs./hd./day.}$$

If he had a set of large framed yearling steers, the adjustment factor would be 1.23. A set of small framed steer calves would have an adjustment factor of .84.

#### Adjustment for Feeding Rumensin

Rumensin is a feed additive for beef cattle that improves feed efficiency by increasing the energy available from a given amount of ration. Daily dry matter intake is reduced approximately 10% when rumensin is fed at the recommended rate, and daily gains are unchanged, resulting in a 10% reduction in feed requirements.

Due to the reduction in dry matter intake and no change in daily gain, some changes must be made in requirement values used to balance the ration. Although the animal is eating 10% less, its daily protein, mineral and salt requirements in actual lbs. are the same because the rate of gain is unchanged. *Thus, the percentages of protein, mineral and salt in ration dry matter given in Tables 1 and 2 in fact sheet 1097 must be 10% higher.* To do this, multiply the percentage requirement by 1.1 and multiply the expected dry matter intake by .9.

The recommended level of feeding rumensin is 30 grams per ton of dry matter or 150 milligrams/head under 600 lb., 200 milligrams from 600 to 800 lb., and 300 milligrams/head/day to those over 800 lb., based on expected daily dry matter intakes.

However, the amounts of supplement to feed/head/day recommended in fact sheets 1204A and 1204B would not be greatly changed. By following those recommendations you are feeding the same amount of supplemental protein and minerals even though rumensin is fed and total ration dry matter intake is reduced.